


EMRO Region

Hydromorphone Consumption (mg/person), 1985 - 2017



Sources: International Narcotics Control Board; The World Bank

Created by: Walther Center in Global Palliative Care & Supportive Oncology (CiGPC), Indiana University Simon Cancer Center, 2020

EMRO Hydromorphone Consumption, (mg/person)

| Year | Hydromorphone |
|------|---------------|
| 1985 | 7.53E-05 |
| 1986 | 0 |
| 1987 | 0 |
| 1988 | 0 |
| 1989 | 0 |
| 1990 | 1.89E-05 |
| 1991 | 3.95E-05 |
| 1992 | 3.60E-05 |
| 1993 | 6.26E-05 |
| 1994 | 7.07E-05 |
| 1995 | 0.0001 |
| 1996 | 2.31E-06 |
| 1997 | 0 |
| 1998 | 3.97E-05 |
| 1999 | 4.74E-05 |
| 2000 | 1.89E-05 |
| 2001 | 6.60E-05 |
| 2002 | 0.0005 |
| 2003 | 5.91E-05 |
| 2004 | 9.25E-05 |
| 2005 | 0.0005 |
| 2006 | 0.0006 |
| 2007 | 0.0003 |
| 2008 | 0.0007 |
| 2009 | 0.0029 |
| 2010 | 0.0029 |
| 2011 | 0.0026 |
| 2012 | 0.0022 |
| 2013 | 0.0035 |
| 2014 | 0.0009 |
| 2015 | 0.0034 |
| 2016 | 0.0105 |
| 2017 | 0.0047 |

Data sources:

Consumption Data – International Narcotics Control Board
Population – The World Bank, 2019 Revision
ME Conversion Factors – WHOCC Centre for Drug Statistics Methodology

Walther Center in Global Palliative Care
& Supportive Oncology
Indiana University
Melvin and Bren Simon Cancer Center